



Clinical Documentation: A Future-Forward Approach

Part 1

Clinical Documentation Intelligence A Guide for Clinicians

Introduction

About the Presenter



Barb Whitlock

Lead Consultant

Healthcare Performance Improvement

Core Objectives

By the end of this session, participants should be able to:

Explain

How documentation shapes clinical signal, risk prediction, throughput, & care variation.

Document

In ways that support artificial intelligence (AI) models, predictive analytics, & care redesign.

Reduce friction

Across care teams by creating clarity, not simply filling gaps.

Documentation is clinical work, not clerical work.



AI Is the Umbrella

These are the parts that read your notes.

Artificial Intelligence



Large Language Models (LLMs)

- Summarize and draft from the narrative record
- Answer questions across long, messy charts
- Only as reliable as the language you write

Natural Language Processing (NLP)

- Extracts terms, findings, & negation from prose
- Feeds coding, quality, & risk adjustment logic
- Ambiguous wording becomes ambiguous data

Both read the same note. Neither can infer what you did not write.



Why Documentation Matters More Now

Documentation is the raw material for care clarity, analytics, risk, quality, & operational performance.

Clinical Signal

Specific, timely documentation helps care teams see trajectory, acuity, response to therapy, & risk.

Operational Signal

The same record informs status, progression, barriers, avoidable days, & transition readiness.

Quality Signal

Documented clinical reality feeds quality measurement, outcomes analysis, & patient safety learning.

Financial Signal

Documentation impacts capture of patient complexity, defensibility, denials, & revenue integrity.

Future-forward documentation is not “more words.” It is more usable clinical evidence.



Two Parallel Data Streams

Most clinicians have never been shown which one their words land in.

Structured

- Problem list & encounter diagnoses
- Orders & medications
- Vitals, flowsheets, I/O, & weights
- Lines, drains, airways
- Scales: MEWS/NEWS, CAM, Braden, Morse
- Labs, imaging, procedures, disposition


Read directly & reliably. Fast, but shallow. It knows a diagnosis exists, not whether it is real, active, or improving.

Unstructured

- HPI & assessment/plan narrative
- Daily progress notes
- Consultant impressions
- Operative & procedure notes
- Nursing & therapy narrative
- Care management & discharge summary

Where the clinical reasoning lives. Rich, but only as good as the language used. Vague prose produces vague inference.

Signal exists where the two streams agree. That agreement is the work.



Four Properties of a Strong Clinical Signal

1

Specificity

What exactly is it?

Weak: "Heart failure"

Strong: "Acute on chronic decompensated HFrEF, EF 25–30%, ischemic etiology"

2

Support

How do you know?

Weak: "Sepsis"

Strong: "Sepsis due to E. coli UTI — T 38.9, HR 118, WBC 18.4, lactate 3.2, MAP 62 requiring 30 mL/kg bolus"

3

Consistency

Does everyone say the same thing?

Weak: "Urosepsis"/"bacteremia without sepsis"

Strong: One reconciled term used by every author, or an explicit, reasoned disagreement

4

Recency

Where is it today, compared to what?


Weak: "Continue diuresis"

Strong: "Net -2.6 L over 48h; O₂ 4 L → 2 L; orthopnea 3 pillows → 1"

From Clinical Note to Operational Insight

Structured & unstructured documentation both become data.


The record is both narrative & data infrastructure.



- Narrative explains clinical reasoning & context.
- Structured data enables measurement, modeling, dashboards, & alerts.
- The feedback loop is only as reliable as the documentation feeding it.

The Four Anchors: 20 Seconds per Problem

Static documentation tells a reader—human or machine—nothing about direction. Direction is what everyone downstream is trying to determine.



Applied: "Dyspnea improved from at rest on admission to exertional beyond 30 feet today; net -1.8 L on IV furosemide 40 mg BID; oxygen weaned 4 L → 2 L."

Timestamp the Inflection Point

An inflection point is the moment the course changed direction. Deterioration models are built to detect this, & validity reviewers are looking for this. If it is not timestamped, neither can find it.

Weak



"Patient required BiPAP overnight."

Strong



"0410 — oxygen requirement escalated 2 L → 6 L with RR 30 and accessory muscle use; ABG 7.29 / 64; BiPAP 14/6 initiated at 0425 with improved work of breathing within 30 minutes."

Four data points. One sentence.

1 | The time 2 | The trigger 3 | The intervention 4 | The interval to response

Make the Chronic Burden Visible

Acute-only documentation makes a complex patient read as a simple one.

The Predictable Results



- An unrealistic expected length of stay
- An unfair observed-to-expected ratio
- Post-acute authorization denied for lack of acuity
- A population health cohort the patient never enters
- A next clinician who inherits half the picture

Document With Status & Stage



- CKD with stage & eGFR · HF with type & EF
- Diabetes with control & named complications
- COPD with severity & home therapy
- Obesity with BMI · cognitive impairment with severity
- Device or support dependence: home O₂, BiPAP, dialysis, feeding tube, ICD

Two rules:

1. Link conditions when a causal relationship exists & you believe it.
2. Do not list what you are not addressing—40 active problems communicates nothing.

Rewriting for Signal

Commonly Written

Rewritten With Signal

What Changes Downstream

"CHF exacerbation. Continue Lasix. Cardiology following."

"Acute on chronic decompensated HFrEF (EF 30%). Baseline NYHA II, now NYHA IV with orthopnea and PND. +8 lb over dry weight. Day 2: net -1.8 L on IV furosemide 40 mg BID, creatinine stable 1.1 → 1.2, O₂ 4 L → 2 L."

- Severity & response reach the risk model
- CM sees a home weight-monitoring need
- UR sees IV titration supporting inpatient status

"Altered mental status. Will monitor."

"Hyperactive delirium (CAM positive 7/14) superimposed on moderate dementia. Precipitants: E. coli UTI and tramadol — tramadol discontinued. Baseline per daughter: oriented, manages own medications."

- Delirium enters fall & deterioration models
- Acute separated from baseline
- Discharge plan can target reversibility

"Diabetes. Continue home meds."

"Type 2 diabetes, uncontrolled hyperglycemia (A1c 9.4%), with stage 3b diabetic CKD (eGFR 38) and peripheral neuropathy. Basal insulin adjusted for renal function."

- Chronic burden visible for risk adjustment
- Linkage supports CKD staging
- Ambulatory team can act on the control gap

High-Value Clinical Descriptors

What clinicians document when they want the record to work harder.



Severity

Mild/moderate/severe; acute, chronic, acute-on-chronic; complication or comorbidity significance



Trajectory

Improving, worsening, fluctuating, refractory, resolved, expected course, discharge readiness




Response

Response to fluids, diuresis, bronchodilators, antibiotics, oxygen, transfusion, procedures, therapy



Risk Context

Frailty, cognition, functional baseline, caregiver support, social risk, clinical vulnerability



Uncertainty

Differential diagnosis, rule-out logic, suspected condition, monitoring plan, why a diagnosis remains possible



Clinical Rationale

Why inpatient care, observation, escalation, discharge delay, specialty consult, or additional testing is needed

Interdisciplinary Documentation Alignment

One record, many readers. Most rework is a translation failure, not a writing failure.

One Record, Many Readers

Reader	What They Need From Your Note	What Breaks When It's Missing
Next Treating Clinician	Current status, trajectory, active plan, contingency if it fails	Duplicated testing, plans reversed at handoff, delayed escalation
CDI	Specificity, clinical indicators, linkage, resolution of uncertainty	Queries, coding delays, unbilled days, validity exposure
Utilization Review	Severity of illness & intensity of service—why this level of care today	Level-of-care mismatch, denials, patient financial harm
Case Management	Baseline & current function, support at home, barriers with dates & owners	Late discharge planning, avoidable days, failed authorization
Ambulatory/PCP	What changed, what is pending, what to follow & when	Dropped follow-up, medication errors, readmission
Quality & Analytics	Comorbidity burden, present-on-admission status, severity	Distorted O/E ratios, misclassified outcomes, unfair reporting

Avoidable Days Are Documented Into Existence—or Out of It



Weak

✕

"Awaiting SNF placement."

Strong

✓

"Medically appropriate for SNF-level care as of 7/14. PT recommends SNF—max assist for transfers, unable to manage stairs. Insurance authorization submitted 7/14 at 1400, pending. Four facilities contacted; two declined for IV antibiotic requirement. Continues IV ceftriaxone q24h through 7/18. Next payer update expected 7/16."

- | **State discharge criteria on day one or two**, not on the day of discharge. Every later note is then measured against a target the whole team can see.
- | **Document the barrier when it appears**, not when it resolves. A barrier documented three days late produces three days no reviewer can defend.

Ambiguity Is a Safety Problem Before It's a Data Problem

Ambiguous Term	Why It Fails	Write Instead
"Urosepsis"	Means sepsis to one reader & a UTI to another	Either "sepsis due to UTI" with indicators, or "UTI without sepsis—no organ dysfunction"
"Altered mental status"	A symptom, not a diagnosis; masks delirium	Name the syndrome & the baseline: delirium, encephalopathy with cause, dementia, intoxication
"Elevated troponin"	A result, not an interpretation; the next clinician must re-adjudicate	State the attribution: type 2 MI from demand ischemia, myocarditis, or CKD without ACS
"History of" an active condition	Reads as resolved & drops the condition from the picture	Reserve for resolved conditions; active chronic conditions get current status & stage
"Possible/likely/probable" at discharge	Uncertainty is fine during the stay, not at the end of it	Confirm, rule out, or state plainly that it is undetermined & what will settle it

When the Record Argues With Itself

A hospitalist, a nephrologist, & an intensivist each describe the same physiology differently. Nobody reconciles it. The record now contains an unresolved argument, & a payor gets to choose which version to believe.

1

Disagree Explicitly, With Reasoning

"I do not believe the troponin elevation represents a type 1 MI: no ischemic ECG changes, no regional wall motion abnormality, and the rise is consistent with demand ischemia in rapid AF."

This is stronger documentation than silence, not weaker.

2

One Author per Problem, per Day

The consultant owns the assessment of their organ system. The attending owns synthesis.

Two simultaneous authoritative assessments of one problem create conflict by default.

3

Resolve on the Day of Resolution

Use the word "resolved" & the date, the moment it is true.

An acute diagnosis carried forward for six days looks either untreated or invented.

Documentation for Predictive Analytics & Risk Modeling

Your documentation now acts on the patient during the stay, not on the claim after it.

From DRG Influence to Risk Influence

Then

Documentation as a Billing Artifact

- One reader, months later
- Which DRG? Which SOI & ROM?
- Will the claim survive review?
- Feedback arrived long after discharge

Getting it wrong costs money after the fact.

Now

Documentation as an Operational Input

- Many readers, continuously
- Deterioration & sepsis alerts; predicted LOS
- Readmission & mortality risk; palliative triggers
- Post-acute authorization; ambulatory rising-risk
- Ambient AI summaries other clinicians read instead of your note

Getting it wrong may change what happens to the patient in the next 12 hours.

Late or thin documentation produces late or thin prediction.
Contradictory documentation produces false alarms the whole unit learns to ignore.



These models are trained
on the record you write.
They cannot see the patient.

They can only see what you said about the
patient, & they cannot tell the difference
between a condition that resolved & a
condition you forgot to remove.



Documentation for Predictive Analytics & Risk Modeling

The shift is from DRG influence only to risk-of-deterioration, LOS prediction, readmission risk, and care redesign.

What Models Need

Specific, timely clinical conditions, severity, trajectory, labs/vitals, treatment intensity, & context

What Teams Need

Operationally useful risk strata that help prioritize work, escalation, & discharge planning

What Leaders Need

Reliable feedback loops for care variation, LOS, denials, quality, & resource planning

Predictive models do not fix poor signal. They amplify the signal they receive.



Avoiding Data Pollution & Documentation Debt

Bad signal can be as harmful as missing signal.



Copy-Forward Noise

Old diagnoses, old status, old plans, stale medication statements



Contradictions

One section says acute, another says chronic or resolved



Non-Actionable Labels

Problem names without severity, treatment linkage, or clinical rationale



Late Documentation

Important facts entered after the window when teams needed them

Clean Signal Principles

- Update the active story daily.
- Resolve contradictions quickly.
- Link diagnoses to evidence & treatment.
- Get critical documentation into the record before decisions are made.

High-Value Descriptors: Greatest Influence per Word Written

Severity & Stage

- EF & HF type
- NYHA
- GOLD & FEV1
- CKD stage with eGFR
- Child-Pugh
- Cancer stage

Functional Status

- ADL dependence
- Assist level for transfers
- Ambulation distance
- Device
- Falls in the past year

Care Intensity

- Frequency of reassessment
- Titratable IV therapy
- Continuous monitoring
- Why today

Acuity & Timing

- Acute/chronic/acute-on-chronic
- Date of onset
- Present on admission vs. hospital-acquired

Cognitive Status

- Documented baseline vs. current
- Delirium screen result & date
- Capacity assessment

Social & Behavioral

- Caregiver availability
- Housing
- Transportation
- Medication cost
- Health literacy

Support Dependence

- O₂ in L/min vs. home baseline
- BiPAP settings
- Pressors
- Dialysis
- Enteral or parenteral nutrition

Nutrition

- BMI
- Percentage & interval of unintentional weight loss
- Malnutrition criteria, provider-affirmed

Goals of Care

- Code status
- Palliative involvement
- Treatment limitations
- Without these, outcome models misread the trajectory



Function, cognition, nutrition, & social barriers are clinical findings. They carry more predictive weight than the primary diagnosis.

Data Pollution: How the Record Suppresses Its Own Accuracy

Documentation that is present, plausible, & wrong—or right but stale. It is the dominant reason predictive tools underperform in real deployments, & it is entirely produced by well-intentioned clinicians under time pressure.

Where It Comes From

- Copy-forward carrying resolved problems & stale exams
- Problem lists that are never pruned
- Templated normal ROS that contradicts the narrative
- Auto-populated data making old text look current
- Placeholder or uncorrected dictation left in a signed note
- Diagnoses entered "to be safe" without indicators
- Conflicting assessments from multiple authors, same day

What It Costs

- **Diluted severity:** When everything is a problem, nothing is prioritized by the model or the next clinician.
- **False positives, then alert fatigue:** Staff dismiss the alert reflexively, including the true ones.
- **Suppressed true risk:** The deteriorating patient is buried in a population that all looks equally sick on paper.
- **Validity denials & audit exposure:** A record that contradicts itself hands a reviewer the contradiction for free.

Six Countermeasures: Leave With One Commitment

1

Copy the Data, Write the Assessment

Rebuild the A/P daily; let the objective section auto-populate.

2

Resolve on the Day of Resolution

Type "resolved" with a date the moment it is true.

3

One Author per Problem, per Day

Consultant owns the organ system; attending owns synthesis.

4

No Placeholders in a Signed Note

Correct dictation errors the same day, before signature.

5

Prune the Problem List

Quarterly in ambulatory; at admission & discharge in acute care or follow your organizations policy

6

Affirm the Team's Findings

Restate dietitian, therapy, & nursing findings in your own assessment.

Case Scenarios

- Heart Failure
- COPD & Asthma
- Frailty & Cognitive Vulnerability

Case Scenario A

Heart Failure: Functional Class & Volume Responsiveness

As Commonly Written

"72-year-old with CHF exacerbation. IV Lasix. Continue diuresis. Cardiology following."

What Is Missing?

- Type & ejection fraction
- Baseline functional class vs. current
- Volume assessment with a reference weight
- Diuretic dose & measured response
- Renal & electrolyte response
- Oxygen requirement & trend
- Precipitant
- Discharge criteria

Rebuilt

"Acute on chronic decompensated HFrEF, EF 25–30%, ischemic etiology. Baseline NYHA II—walks two blocks without stopping; presented NYHA IV with three-pillow orthopnea and PND. Volume overload: 9 lb above dry weight of 78 kg, JVP 12 cm, 2+ pitting edema to the knees. Precipitant: dietary sodium plus five missed doses of torsemide—patient reports a copay barrier.

Hospital day 2: net –2.6 L, weight down 4 lb, creatinine 1.2 → 1.3 (acceptable, continuing), K 3.4 repleted, oxygen 3 L → room air at rest, dyspnea now only with ambulation beyond 50 feet. Continue IV furosemide 40 mg BID; convert to oral at dry weight.

Discharge criteria: room air at rest and with ambulation, 24 hours of stable weight on oral diuretic, medication-assistance enrollment completed, home health with weight monitoring arranged."

Case Scenario A

Heart Failure: What Changed Downstream

Risk Adjustment

Severity & present-on-admission status reach the expected-mortality calculation

Readmission Model

Receives two of its strongest predictors: functional class & a medication-cost adherence barrier

Case Management

Can act on the copay barrier on day two instead of discovering it after readmission

Utilization Review

Sees active IV titration with measured response: the intensity of service supporting inpatient status

PCP

Inherits a dry weight, a target regimen, & a reason the patient decompensated

Teaching Points

- Always name type & EF. "CHF" alone is the highest-volume specificity gap in most organizations.
- Express functional class in distance or activity. "Two blocks" is data; "limited" is not.
- Document diuretic response with numbers: net balance, weight, creatinine, oxygen.
- Name the precipitant—the single most actionable element for preventing the next admission.
- State the dry weight. Nearly every post-discharge intervention depends on it, & it is usually absent.

Case Scenario B

COPD & Asthma: Disease Severity & Home Therapy Needs

As Commonly Written

"COPD exacerbation. Steroids, nebs, continue current management."

What Is Missing?

- Baseline severity & exacerbation history
- Home oxygen & noninvasive ventilation baseline
- Degree of deviation from that baseline
- Escalation moment
- Trigger
- Inhaler technique & adherence
- Discharge criteria referenced to home therapy

Rebuilt

"Acute exacerbation of severe COPD, GOLD group E—FEV1 38% predicted (3/2025), two exacerbations and one admission in the past twelve months. Chronic home therapy: oxygen 2 L continuous and nocturnal BiPAP 14/6.

Presented with RR 30, accessory muscle use, pH 7.31 and pCO₂ 62 against a baseline pCO₂ of approximately 50; BiPAP initiated in the ED at 2140.

Hospital day 2: off BiPAP for 14 hours, requiring 3 L—still above home baseline—RR 20, speaking in full sentences, ambulated 40 feet with SpO₂ nadir 88% on 3 L. Trigger: viral URI; smoking cessation maintained two years. RT assessment shows poor inhaler technique—re-education completed, spacer ordered.

Discharge criteria: back to home oxygen of 2 L with ambulation SpO₂ above 88%, off daytime BiPAP, steroid course plan documented, pulmonary follow-up within 7 days."

Case Scenario B

COPD & Asthma: Teaching Points

- 1 | Anchor Severity to Something Objective**
FEV1, GOLD group, exacerbation & admission history, or prior intubation—not the word "exacerbation"
 - 2 | Document the Home Baseline First**
Without it, no reader can tell how far from normal the patient actually is
 - 3 | Document Deviation, Not Just Abnormality**
"3 liters" reads as normal until you say the patient lives on two.
 - 4 | Capture the Escalation Moment With a Time**
A note that says "needed BiPAP" loses the entire inflection point
 - 5 | Technique & Adherence Are Clinical Findings**
They belong in your assessment, not only in the respiratory therapy note.
- Downstream:** The timestamped escalation is what both the deterioration model & any later medical necessity review need; technique & follow-up interval feed readmission prediction.

Case Scenario C

Frailty & Cognitive Vulnerability

As Commonly Written

"Elderly, deconditioned, confused at times. PT to evaluate. Likely needs rehab."

What Is Missing?

- Baseline for both function & cognition
- Whether the confusion is acute or chronic
- Frailty & fall history
- Nutritional status
- Precipitants of any acute change
- Caregiver picture
- Trajectory of function during the stay

Rebuilt

"Frailty: Clinical Frailty Scale 6 at baseline—per daughter, requires assistance with bathing and dressing, uses a walker indoors, stopped cooking six months ago.

Nutrition: unintentional weight loss of 7 kg over six months (9% of body weight); dietitian documents criteria for moderate malnutrition and I agree—treating with oral supplementation and monitoring intake.

Cognition: moderate dementia at baseline (MoCA 16, 4/2025) with superimposed hypoactive delirium—CAM positive on admission 7/14; precipitants were E. coli UTI and oxybutynin, which has been discontinued. 7/16: CAM negative, oriented to person and place, family confirms return to baseline.

Falls: two in the past six months, one with a wrist fracture; Morse score 65. Function: maximum assist for transfers on admission, now minimum assist; PT recommends SNF-level rehabilitation."

Case Scenario C

Frailty & Cognition: Why This One Matters Most

Among the Strongest Predictors

Frailty, malnutrition, & functional dependence predict mortality, LOS, & readmission. None of them exist in the data if only the therapist writes them down.

Delirium Is Reversible; Dementia Is Not

Separating them keeps an acute, treatable problem from being recorded as a permanent baseline that follows the patient into every future encounter.

Baseline Makes Improvement Measurable

It is also exactly what post-acute authorization requires. Without it, the SNF request has nothing to stand on.

The Caregiver Detail Is the Discharge Plan

It is the difference between a safe discharge & a bounce-back.

The rule that closes the gap: The provider must affirm dietitian & therapy findings in their own assessment for those findings to enter the data at all.

Assessment Knowledge Check

1

A diagnosis appears in your narrative but was never added to the problem list. Most likely consequence?

It may be invisible to structured logic—cohort identification, rules-based risk stratification, quality abstraction.

2

Why does an unpruned problem list degrade model performance?

It dilutes severity & generates false positives, driving alert fatigue & masking genuinely deteriorating patients.

3

A patient waits three days for placement. What protects those days?

Daily documentation of the specific barrier, action taken, date & time, owner, & next milestone.

4

Which is the strongest documentation of therapeutic response?

"Net -2.6 L over 48 hours; weight down 4 lb; oxygen 4 L → room air; orthopnea 3 pillows → 1."—state, comparison, intervention, measured response.

5

A dietitian documents severe malnutrition criteria. What must happen next?

The provider must affirm the diagnosis in their own assessment & document treatment.

6

Which term should leave clinical documentation entirely?

"Urosepsis." It means sepsis to one reader & a UTI to another. Say which one you mean, with indicators.

Takeaway

The Documentation Intelligence Checklist

The entire session, compressed into eight habits.

- 1 | Name It Precisely**
Type, acuity, severity, stage, site, etiology
- 2 | Give a Reference Point**
Today vs. yesterday, admission, or stated baseline
- 3 | Timestamp the Inflection**
When the course changed, what triggered it, what you did
- 4 | Resolve & Reconcile**
"Resolved" with a date; explicit, reasoned disagreement
- 5 | Show Your Evidence**
Clinical indicators in the same note as the diagnosis
- 6 | Show the Response**
The intervention & its measured effect—including none
- 7 | Make the Burden Visible**
Chronic conditions being addressed—and only those
- 8 | Write for the Next Reader**
Would they know the direction of travel?

In Closing...

The future of documentation is clarity that clinicians trust & systems can use.

Commitment Prompts:

- What is one documentation phrase we will standardize?
- Where is our highest-friction documentation handoff?
- Which team needs a better feedback loop?
- What can we remove from documentation that creates noise?

Clinical Documentation Intelligence Means:

The right clinical facts are present, specific, timely, consistent, & usable enough to support care, quality, operations, analytics, & audit defense.

Write the record you would want to inherit.



Next in the Series

Defensible Documentation: Protecting Clinical Decisions

Real-world strategies for defending medical decision making, not just DRGs.



Part 1 taught how to make documentation useful. Part 2 teaches how to make it defensible. Every contradiction, ambiguity, & copied paragraph we discussed today is what a reviewer looks for first.

Contact

Forvis Mazars

Barb Whitlock

Lead Consultant

Healthcare Performance Improvement

barb.whitlock@us.forvismazars.com

The information set forth in this presentation contains the analysis and conclusions of the author(s) based upon his/her/their research and analysis of industry information and legal authorities. Such analysis and conclusions should not be deemed opinions or conclusions by Forvis Mazars or the author(s) as to any individual situation as situations are fact-specific. The reader should perform their own analysis and form their own conclusions regarding any specific situation. Further, the author(s)' conclusions may be revised without notice with or without changes in industry information and legal authorities.

© 2026 Forvis Mazars, LLP. All rights reserved.